

**Final
Environmental Impact Statement/Overseas Environmental Impact Statement
Atlantic Fleet Training and Testing**

TABLE OF CONTENTS

3.1	Air Quality	3.1-1
3.1.1	Introduction	3.1-1
3.1.1.1	Air Quality Standards	3.1-2
3.1.1.2	General Conformity Evaluation	3.1-5
3.1.1.3	Approach to Analysis	3.1-7
3.1.1.4	Emission Estimates	3.1-9
3.1.1.5	Climate Change	3.1-11
3.1.1.6	Other Compliance Considerations, Requirements, and Practices	3.1-12
3.1.2	Affected Environment.....	3.1-13
3.1.2.1	General Background	3.1-13
3.1.2.2	Sensitive Receptors.....	3.1-13
3.1.2.3	Existing Air Quality.....	3.1-20
3.1.3	Environmental Consequences	3.1-23
3.1.3.1	Impacts from Air Emissions Under Alternative 1.....	3.1-24
3.1.3.2	Impacts from Air Emissions under Alternative 2	3.1-26
3.1.3.3	Impacts from Air Pollutants under the No Action Alternative	3.1-29
3.1.3.4	Greenhouse Gases and Climate Change.....	3.1-29
3.1.4	Summary of Potential Impacts on Air Quality	3.1-29
3.1.4.1	Combined Impacts of All Stressors under Alternative 1.....	3.1-30
3.1.4.2	Combined Impacts of All Stressors under Alternative 2.....	3.1-30
3.1.4.3	Combined Impacts of All Stressors under the No Action Alternative	3.1-30

List of Figures

Figure 3.1-1: Applicable Nonattainment and Maintenance Areas in USEPA Region 1 and 2.....	3.1-14
Figure 3.1-2: Applicable Nonattainment and Maintenance Areas in USEPA Region 3.....	3.1-15
Figure 3.1-3: Applicable Nonattainment and Maintenance Areas in USEPA Region 4.....	3.1-16
Figure 3.1-4: Applicable Nonattainment and Maintenance Areas in USEPA Region 6.....	3.1-17

List of Tables

Table 3.1-1: National Ambient Air Quality Standards.....	3.1-3
Table 3.1-2: <i>De Minimis</i> Thresholds for Conformity Determinations.....	3.1-7
Table 3.1-3: Nonattainment and Maintenance Areas Adjacent to Study Area	3.1-21
Table 3.1-4: Pierside and Coastal Activity Locations and Their Area’s Attainment Status	3.1-22
Table 3.1-5: Estimated Annual Air Pollutant Emissions from Activities Occurring within the AFTT Study Area, Alternative 1	3.1-24
Table 3.1-6: Estimated Annual Air Pollutant Emissions from Activities Occurring in State Waters in the Jacksonville, Florida Area, Alternative 1	3.1-25
Table 3.1-7: Estimated Annual Air Pollutant Emissions from Activities Occurring within the AFTT Study Area, Alternative 2	3.1-27
Table 3.1-8: Estimated Annual Air Pollutant Emissions from Activities Occurring within 3 NM of shore in the Jacksonville, Florida Area, Alternative 2.....	3.1-28
Table 3.1-9: Total Annual Greenhouse Gas Emissions from All Study Area Training and Testing Activities in Metric Tons per Year	3.1-29

3.1 AIR QUALITY

AIR QUALITY SYNOPSIS

The United States Department of the Navy considered all potential stressors that air quality could be exposed to from the Proposed Action. The following conclusions have been reached for the Preferred Alternative (Alternative 1):

- Criteria Air Pollutants: The emission of criteria pollutants resulting from activities in the Study Area would not cause a violation or contribute to an ongoing violation of the National Ambient Air Quality Standards.

3.1.1 INTRODUCTION

Air pollution is a threat to human health and also damages the environment (U.S. Environmental Protection Agency, 2007). Air pollution damages trees, crops, other plants, lakes, and animals. In addition to damaging the natural environment, air pollution damages the exteriors of buildings, monuments, and statues. It creates haze or smog that reduces visibility in national parks and cities and interferes with aviation. To improve air quality and reduce air pollution, Congress passed the Clean Air Act and its amendments in 1970 and 1990, which set regulatory limits on air pollutants and help to ensure basic health and environmental protection from air pollution.

Air quality is defined by ambient concentrations of specific air pollutants – pollutants the U.S. Environmental Protection Agency (USEPA) determined may affect the health or welfare of the public. The six major pollutants of concern are called “criteria pollutants”: carbon monoxide, sulfur dioxide, nitrogen dioxide, ozone, particulate matter (dust particles less than or equal to 10 microns in diameter and fine particulate matter less than or equal to 2.5 microns in diameter), and lead. The Clean Air Act required that the USEPA establish National Ambient Air Quality Standards for these criteria pollutants. These standards set specific concentration limits for criteria pollutants in the outdoor air. The concentration limits were developed because the criteria pollutants are common in outdoor air, considered harmful to public health and the environment, and come from numerous and diverse sources. The concentration limits are designed to aid in protecting public health and the environment. Areas with air pollution problems typically have one or more criteria pollutants consistently present at levels that exceed the National Ambient Air Quality Standards. These areas are designated as nonattainment for the standards.

Criteria air pollutants are classified as either primary or secondary pollutants based on how they are formed in the atmosphere. Primary air pollutants are emitted directly into the atmosphere from the source of the pollutant and retain their chemical form. Examples of primary pollutants are the smoke produced by burning wood and volatile organic compounds emitted by industrial solvents. Secondary air pollutants are those formed through atmospheric chemical reactions that usually involve primary air pollutants (or pollutant precursors) and normal constituents of the atmosphere. Ozone, a major component of photochemical smog, is a secondary air pollutant. Ozone precursors fall into two broad groups of chemicals: nitrogen oxides and volatile organic compounds. Nitrogen oxides consists of nitric oxide and nitrogen dioxide.

Finally, some criteria air pollutants are a combination of primary and secondary pollutants. Particulate matter less than or equal to 10 microns in diameter and particulate matter less than or equal to 2.5 microns in diameter are generated as primary pollutants by various mechanical processes (e.g., abrasion, erosion, mixing, or atomization) or combustion processes. They are generated as secondary pollutants through chemical reactions or through the condensation of gaseous pollutants into fine aerosols.

In addition to the six criteria pollutants, the USEPA currently designates 187 substances as hazardous air pollutants under the federal Clean Air Act. Hazardous air pollutants are air pollutants known or suspected to cause cancer or other serious health effects, or adverse environmental and ecological effects (U.S. Environmental Protection Agency, 2016a). National Ambient Air Quality Standards are not established for these pollutants; however, the USEPA developed rules that limit emissions of hazardous air pollutants from specific industrial sources. These emissions control standards are known as “maximum achievable control technologies” and “generally achievable control technologies.” They are intended to achieve the maximum degree of reduction in emissions of the hazardous air pollutants, taking into consideration the cost of emissions control, non-air quality health and environmental impacts, and energy requirements. These emissions are typically one or more orders of magnitude smaller than concurrent emissions of criteria air pollutants, and only become a concern when large amounts of fuel, explosives, or other materials are consumed during a single activity or in one location. Hazardous air pollutants are analyzed qualitatively in relation to the prevalence of the sources emitting these pollutants during training and testing activities. Mobile sources operating as a result of the Proposed Action would be functioning intermittently over a large area and would produce negligible ambient hazardous air pollutants, predominantly in areas not routinely accessed by the general public. For these reasons, hazardous air pollutants are not further evaluated in the analysis. Air pollutant emissions are reported as the rate (by weight or volume) at which specific compounds are emitted into the atmosphere by a source. Most air pollutant emissions are expressed as a rate (e.g., pounds per hour, pounds per day, or tons per year). Typical units for emission factors for a source or source activity are pounds per thousand gallons of fuel burned, pounds per ton of material processed, and grams per vehicle-mile of travel.

Ambient air quality is reported as the atmospheric concentrations of specific air pollutants at a particular time and location. The units of measurement are expressed as a mass per unit volume (e.g., micrograms per cubic meter [$\mu\text{g}/\text{m}^3$] of air) or as a volume fraction (e.g., parts per million [ppm] by volume). The ambient air pollutant concentrations measured at a particular location are determined by the pollutant emissions rate, local meteorology, and atmospheric chemistry. Wind speed and direction, the vertical temperature gradient of the atmosphere, and precipitation patterns affect the dispersal, dilution, and removal of air pollutant emissions from the atmosphere.

3.1.1.1 Air Quality Standards

National Ambient Air Quality Standards for criteria pollutants are set forth in Table 3.1-1. Areas that exceed a standard are designated as “nonattainment” for that pollutant, while areas that are in compliance with a standard are in “attainment” for that pollutant. An area may be nonattainment for some pollutants and attainment for others simultaneously.

Table 3.1-1: National Ambient Air Quality Standards

<i>Pollutant</i>		<i>Primary/ Secondary</i>	<i>Averaging Time</i>	<i>Level</i>	<i>Form</i>
Carbon monoxide		primary	8 hours	9 ppm	Not to be exceeded more than once per year
			1 hour	35 ppm	
Lead		primary and secondary	Rolling 3-month period	0.15 µg/m ³ ⁽¹⁾	Not to be exceeded
Nitrogen dioxide		primary	1 hour	100 parts per billion (ppb)	98th percentile of 1-hour daily maximum concentrations, averaged over 3 years
		primary and secondary	1 year	53 ppb ⁽²⁾	Annual mean
Ozone		primary and secondary	8 hours	0.070 ppm ⁽³⁾	Annual fourth-highest daily maximum 8-hour concentration, averaged over 3 years
Particle pollution (particulate matter)	particulate matter less than or equal to 2.5 microns in diameter	primary	1 year	12.0 µg/m ³	Annual mean, averaged over 3 years
		secondary	1 year	15.0 µg/m ³	Annual mean, averaged over 3 years
		primary and secondary	24 hours	35 µg/m ³	98th percentile, averaged over 3 years
	particulate matter less than or equal to 10 microns in diameter	primary and secondary	24 hours	150 µg/m ³	Not to be exceeded more than once per year on average over 3 years

Table 3.1-1: National Ambient Air Quality Standards (continued)

<i>Pollutant</i>	<i>Primary/ Secondary</i>	<i>Averaging Time</i>	<i>Level</i>	<i>Form</i>
Sulfur dioxide	primary	1 hour	75 ppb ⁽⁴⁾	99th percentile of 1-hour daily maximum concentrations, averaged over 3 years
	secondary	3 hours	0.5 ppm	Not to be exceeded more than once per year

⁽¹⁾In areas designated nonattainment for the lead standards prior to the promulgation of the current (2008) standards, and for which implementation plans to attain or maintain the current (2008) standards have not been submitted and approved, the previous standards (1.5 µg/m³ as a calendar quarter average) also remain in effect.

⁽²⁾The level of the annual nitrogen dioxide standard is 0.053 ppm. It is shown here in terms of ppb for the purposes of clearer comparison to the 1-hour standard level.

⁽³⁾Final rule signed October 1, 2015, and effective December 28, 2015. The previous (2008) ozone standards additionally remain in effect in some areas. Revocation of the previous (2008) ozone standards and transitioning to the current (2015) standards will be addressed in the implementation rule for the current standards.

⁽⁴⁾The previous sulfur dioxide standards (0.14 ppm 24-hour and 0.03 ppm annual) will additionally remain in effect in certain areas: (1) any area for which it is not yet 1 year since the effective date of designation under the current (2010) standards, and (2) any area for which implementation plans providing for attainment of the current (2010) standard have not been submitted and approved and which is designated nonattainment under the previous sulfur dioxide standards or is not meeting the requirements of a State Implementation Plan call under the previous sulfur dioxide standards (40 Code of Federal Regulations [CFR] 50.4(3)). A State Implementation Plan call is a USEPA action requiring a state to resubmit all or part of its State Implementation Plan to demonstrate attainment of the require National Ambient Air Quality Standards.

Source: (U.S. Environmental Protection Agency, 2016b), last updated January 7, 2016.

Notes: µg/m³ = micrograms per cubic meter; ppb = parts per billion; ppm = parts per million

States, through their air quality management agencies, are required to prepare and implement State Implementation Plans for nonattainment areas, which demonstrate how the area will meet the National Ambient Air Quality Standards. Areas classified as attainment, after being designated as nonattainment, may be reclassified as maintenance areas subject to maintenance plans showing how the area will continue to meet federal air quality standards. Nonattainment areas for some criteria pollutants are further classified, depending on the severity of their air quality problem, to facilitate their management:

- ozone – marginal, moderate, serious, severe, and extreme
- carbon monoxide – moderate and serious
- particulate matter – moderate and serious

The USEPA delegates the regulation of air quality to the state once the state has an approved State Implementation Plan. If the state fails to develop an adequate plan to achieve and maintain the National Ambient Air Quality Standards or a State Implementation Plan revision is not approved by the USEPA, federal agencies must comply with the Federal Implementation Plan. States may also choose to adopt the Federal Implementation Plan as an alternative to developing their own State Implementation Plan. States may establish air quality standards more stringent than the National Ambient Air Quality Standards, however they are prohibited from imposing more stringent conformity requirements unless the requirements apply equally to non-federal activities.

The Atlantic Fleet Training and Testing (AFTT) Study Area is offshore of a number of states, and some elements of the Proposed Action occur within or over state waters. State waters extend from the shoreline to 3 nautical miles (NM) from Maine to the east coast of Florida, Alabama, Mississippi, Louisiana, and to 9 NM for the west coast of Florida and Texas. A coastal state exercises sovereignty over its territorial sea, the air space above it, and the seabed and subsoil beneath it. Some activities occur in state waters and primarily involve the use of small boats as is the case with inshore training on state waters. These activities occur in a variety of locations such as Narragansett Bay, the lower Chesapeake Bay, the James and York Rivers, Kings Bay, Cooper River, St. Johns River, and St. Andrew Bay. However, most of the Study Area is substantially offshore, beyond state boundaries where attainment status is unclassified and Clean Air Act National Ambient Air Quality Standards do not apply. There may be seasonal or other temporal fluctuations in wind direction, and during these periods, air quality in adjacent onshore areas may be affected by releases of air pollutants from mobile sources within the Study Area. Impacts at a scale that would produce demonstrable air quality impacts would typically be the result of heavy marine traffic in areas such as large ports but military activity could incrementally impact these areas. Therefore, National Ambient Air Quality Standards attainment status of adjacent onshore areas is considered in determining whether appropriate controls for air pollution sources in the adjacent offshore state waters is warranted.

In addition to coastal states, training may occur in areas proximate to U.S. territories; specifically, the U.S. Virgin Islands, Puerto Rico, Culebra and Vieques. Territorial seas for the U.S. Virgin Islands are within 3 NM of the islands. For Puerto Rico and its neighboring islands Culebra and Vieques, the territorial seas lie within 9 NM of the coast. Air quality is typically very good at all of these locations, which are heavily influenced by tradewinds and air currents induced by convection.

3.1.1.2 General Conformity Evaluation

Federal actions are required to conform with the approved State Implementation Plan for those areas of the United States designated as nonattainment or maintenance areas for any criteria air pollutant under the Clean Air Act (40 CFR parts 51 and 93). The purpose of the General Conformity Rule is to ensure that applicable federal actions, such as the Proposed Action evaluated in this Environmental Impact Statement (EIS)/Overseas Environmental Impact Statement (OEIS), would not cause or contribute to a violation of an air quality standard and that the Proposed Action would not adversely affect the attainment and maintenance of National Ambient Air Quality Standards. A conformity evaluation must be completed for every applicable Navy action that generates emissions in non-attainment or maintenance areas to determine and document whether a proposed action complies with the General Conformity Rule.

In keeping with Navy guidance, the first step in evaluating the Navy action under General Conformity is to define the action, which is to conduct training activities and research, development, testing, and evaluation activities in the AFTT Study Area. This action will utilize numerous mobile sources, including aircraft, small boats, and ships, and will involve the expenditure of munitions. A detailed explanation of the proposed action can be found in Chapter 2 of this document. A significant portion of the Study Area activities would occur well offshore, but there are nearshore areas where activities will take place. One nearshore area that is located in a nonattainment area is a portion of Nassau County, which is part of the Jacksonville (Florida)-Brunswick (Georgia) Interstate Air Quality Control Region and lies adjacent to Duval County. Because a portion of the proposed Federal action will occur near a nonattainment area and the action involves the combustion of fossil fuel, an assessment has to be made as to whether or not the action is considered exempt or presumed to conform under the General Conformity Regulations.

Exempt actions are very specifically defined by USEPA, and the list of exempt actions is available at 40 CFR 93.153. The action's non-exempt direct and indirect emissions are then calculated to determine the *de minimis* emission levels for the applicable pollutants.

The total direct and indirect emissions is defined as the net emissions increase caused by the action considering all the emission increases and decreases that are projected to occur. The portion of emissions that are exempt or presumed to conform are not included in the total of direct and indirect emissions. The total direct and indirect emissions calculation considers all non-exempt emission increases and decreases, must be reasonably foreseeable at the time that the conformity evaluation is conducted, and the emissions considered in the calculation are within the agency's program responsibility and control (U.S. Environmental Protection Agency, 2017a).

If the total direct and indirect emissions do not exceed *de minimis* thresholds defined in the regulation, then a General Conformity Determination is not required. The *de minimis* thresholds are presented in Table 3.1-2. If these emissions equal or exceed the *de minimis* threshold values, a formal Conformity Determination must be prepared to demonstrate conformity with the USEPA-approved State Implementation Plan. If the total direct and indirect emissions do not exceed *de minimis* thresholds, then a General Conformity Determination is not required. If these emissions equal or exceed the *de minimis* threshold values, a formal Conformity Determination must be prepared to demonstrate conformity with the approved State Implementation Plan.

The Navy Guidance for Compliance with the Clean Air Act General Conformity Rule section 4.1, states that a Record of Non-Applicability must be prepared if the proposed action is subject to the Conformity Rule, but is exempt because it fits within one of the exemption categories listed under 40 CFR 93B, because the action's projected emissions are below the *de minimis* conformity applicability threshold values, or is presumed to conform (U.S. Department of the Navy, 2013).

The Clean Air Act sets out specific requirements for a group of northeast states that make up the Ozone Transport Region. States in this region are required to submit a State Implementation Plan and install a certain level of controls for the pollutants that form ozone, even if they meet the ozone standards. The Ozone Transport Region includes Maine, New Hampshire, Vermont, Massachusetts, Rhode Island, Connecticut, New York, New Jersey, Pennsylvania, Delaware, Maryland, and the Washington, D.C. Metropolitan Statistical Area, including the northern Virginia suburbs (Ozone Transport Commission, 2017). The Ozone Transport Region is an area subjected to poor air quality in the warm summer months resulting from ozone pollution. Contributing to the problem are local sources of air pollution as well as air pollution transported hundreds of miles from distant sources in and outside of the Ozone Transport Region. Transport most frequently originates in the Midwest and the Ohio River Valley. Note that *de minimis* levels for ozone precursors may be lower where nonattainment is a serious issue in the Ozone Transport Region. It should be noted that not all of the Ozone Transport Region has issues with attainment of the National Ambient Air Quality Standards. For example, the entire state of Rhode Island is in attainment for all criteria pollutants. However, its geographical location, as with all of the Ozone Transport Region, makes it potentially susceptible to pollutant emission incursions from other areas of the country.

Table 3.1-2: De Minimis Thresholds for Conformity Determinations

<i>Pollutant</i>	<i>Nonattainment or Maintenance Area Type</i>	<i>de Minimis Threshold (TPY)</i>
Ozone (VOC or NO _x)	Serious nonattainment	50
	Severe nonattainment	25
	Extreme nonattainment	10
	Other areas outside an Ozone Transport Region	100
Ozone (NO _x)	Marginal and moderate nonattainment inside an Ozone Transport Region	100
	Maintenance	100
Ozone (VOC)	Marginal and moderate nonattainment inside an Ozone Transport Region	50
	Maintenance within an Ozone Transport Region	50
	Maintenance outside an Ozone Transport Region	100
CO, SO ₂ and NO ₂	All nonattainment and maintenance	100
PM ₁₀	Serious nonattainment	70
	Moderate nonattainment and maintenance	100
PM _{2.5}	All nonattainment and maintenance	100
Lead	All nonattainment and maintenance	25

Source: 40 CFR part 93B

Notes: CO: carbon monoxide; NO_x: nitrogen oxides; NO₂: nitrogen dioxide; PM₁₀: particulate matter ≤ 10 microns in diameter; PM_{2.5}: particulate matter ≤ 2.5 microns in diameter; SO₂: sulfur dioxide; SO_x: sulfur oxides; TPY: tons per year; VOC: volatile organic compound

3.1.1.2.1 Conformity Analysis in Nonattainment and Maintenance Areas

Certain Navy training and testing activities take place within nonattainment and maintenance areas. These nonattainment and maintenance areas are identified by their air quality designated areas (an area designated by the federal government where communities share a common air pollution problem). Several designated areas were identified as relevant to AFTT EIS/OEIS training or testing activities and are further discussed in Section 3.1.2.3, Existing Air Quality.

3.1.1.3 Approach to Analysis

Boundaries of Analysis

The air quality impact evaluation requires two separate analyses. Impacts of air pollutants emitted by Navy training and testing in the Atlantic Ocean, state waters, bays and inshore locations are assessed under National Environmental Policy Act (NEPA). Impacts of air pollutants emitted by Navy training and testing activities outside territorial waters are evaluated as required under Executive Order 12114.

Air pollutants emitted more than 3,000 feet (ft.) above ground level are considered to be above the atmospheric inversion layer and, therefore, do not affect ground level air quality (U.S. Environmental Protection Agency, 2009a). These emissions thus do not affect the concentrations of criteria air pollutants in the lower atmosphere, which are measured at ground level monitoring stations, and upon which federal, state, and local regulatory decisions are based. For the analysis of the effects on global climate change, however, all emissions of greenhouse gases from aircraft and vessels participating in training and testing activities, as well as targets and munitions expended, are applicable regardless of altitude (Chapter 4, Cumulative Impacts). However, because activities above 3,000 ft. for individual aircraft activities are not specifically documented, it would be impossible to analyze with any accuracy the greenhouse gases associated with testing and training activity flights above 3,000 ft. For this reason,

the greenhouse gas emissions that are assessed should be understood to represent only a portion of the total emissions from aircraft flight activities.

Analysis of health-based air quality impacts under NEPA and Executive Order 12114 includes estimates of criteria air pollutants for all training and testing activities where aircraft, missiles, or targets operate at or below the aforementioned inversion layer or that involve vessels in U.S. territorial seas. The analysis of health-based air quality impacts under Executive Order 12114 includes emissions estimates of only those training and testing activities in which aircraft, missiles, or targets operate at or below 3,000 ft. above ground level, or that involve vessels outside of U.S. territorial seas. While there are a number of pierside locations associated with testing and training activities, emissions that may be generated at these locations are excluded from this analysis because they have already been previously analyzed in other NEPA documentation.

Emission Sources

Criteria air pollutants are generated by the combustion of fuel by surface vessels and by fixed-wing and rotary-wing aircraft. These mobile sources are the primary emitters of air pollution associated with testing and training activities. The emissions from these mobile sources are a function of combustion of fuel and emissions are estimated using information provided by the Navy and other reputable, sanctioned sources. Emissions are also generated by the combustion of explosives and propellants in various types of munitions. Propellants used to fire small-, medium-, and large-caliber projectiles generate criteria pollutants when detonated. Non-explosive practice munitions contain spotting charges and propellants that generate criteria air pollutants when they function. Powered targets require fuel, generating criteria air pollutants during their operation, and towed targets generate criteria air pollutants secondarily because another aircraft or vessel is required to provide power. Stationary targets may generate criteria air pollutants if all or portions of the item burn in a high-order detonation. Chaff cartridges used by ships and aircraft are launched by an explosive charge that generates small quantities of criteria air pollutants. Countermeasure flares, parachute flares, and smoke floats are designed to burn for a prescribed period, emitting criteria pollutants in the process.

The primary emissions from many munition types are carbon dioxide, carbon monoxide, and particulate matter; hazardous air pollutants are emitted at low levels (U.S. Environmental Protection Agency, 2009a).

Electronic warfare countermeasures generate emissions of chaff, a form of particulate not regulated under the federal Clean Air Act as a criteria air pollutant because virtually all radio frequency chaff is 10 to 100 times larger than particulate matter less than or equal to 10 microns in diameter (PM₁₀) (U.S. Department of the Navy, 1999). The types of training and testing that produce these other emissions may take place throughout the Study Area, but occur primarily within special use airspace. Chaff emissions during training and testing primarily occur 3 NM or more from shore and at altitudes over 3,000 ft. (above the mixing layer). Chaff released over the ocean would disperse in the atmosphere and then settle onto the ocean surface.

A study at Naval Air Station Fallon found that the release of 50,000 cartridges of chaff per year over 10,000 square miles (m²) would result in an annual average concentration of 0.018 µg/m³ for regulated particulate matter. This is far below the National Ambient Air Quality Standards. Similar predictions were made for St. Mary's County, Maryland (on the Chesapeake Bay), where chaff releases contribute no more than 0.008 percent of total particulate matter emissions (Arfsten et al., 2001). Therefore, chaff is not further evaluated as an air quality stressor in this EIS/OEIS.

3.1.1.3.1 Analysis Framework

Emissions sources and the approach used to estimate emissions under Alternative 1 and Alternative 2 for the air quality analysis are based, wherever possible, on information from Navy subject matter experts and established training and testing requirements. These data were used to estimate the numbers and types of aircraft, surface ships and vessels, submarines, and munitions (i.e., potential sources of air emissions) that would be involved in training and testing activities under each alternative. Emissions were assessed to identify any possibility for the magnitude of Proposed Action emissions to result in a violation of one or more National Ambient Air Quality Standards. The pollutants for which calculations are made include exhaust total hydrocarbons, carbon monoxide, nitrogen oxides, particulate matter, carbon dioxide, and sulfur dioxide and carbon dioxide.

The NEPA analysis includes a separate section for a Clean Air Act General Conformity Applicability Analysis to support a determination pursuant to the General Conformity Rule (40 CFR part 93B). This analysis focuses on training and testing activities that could impact nonattainment or maintenance areas within the region of influence. As noted above, the Study Area lies partly within or adjacent to some air quality designated areas. To evaluate whether or not the General Conformity Rule applies, air pollutant emissions associated with the Proposed Action within the applicable designated nonattainment or maintenance areas are estimated, based on the distribution of mobile source activity in state waters and mobile source activity beyond state waters. The proposed training and testing activities within this portion of the Study Area are then compared to the General Conformity Rule *de minimis* thresholds.

3.1.1.4 Emission Estimates

3.1.1.4.1 Aircraft Activities

To estimate aircraft emissions, the operating modes, number of hours of operation, and type of engine for each type of aircraft were evaluated.

Emissions associated with airfield or air station operations ashore are analyzed within the home basing environmental planning process (e.g., environmental impact statements or environmental assessments for (1) *Introduction of F/A-18 E/F (Super Hornet) Aircraft to the East Coast of the United States* (U.S. Department of the Navy, 2003); (2) *Supplemental Environmental Impact Statement for the Introduction of the P-8A Multi-Mission Aircraft into the U.S. Navy Fleet* (U.S. Department of the Navy, 2014); (3) *Transition of E-2C Hawkeye to E-2D Advanced Hawkeye at Naval Station Norfolk, Virginia, and Naval Base Ventura County Point Mugu, California* (U.S. Department of the Navy, 2009), and (4) *F-35B East Coast Basing Environmental Impact Statement* (U.S. Department of the Navy, 2010). All fixed-wing aircraft are assumed to travel to and from training and testing ranges at or above 3,000 ft. above mean sea level and, therefore, their transits to and from the ranges do not affect surface air quality. Air combat maneuvers and air-to-air missile exercises are primarily conducted at altitudes well in excess of 3,000 ft. above mean sea level and, therefore, are not included in the estimated emissions of criteria air pollutants. Activities or portions of those training or testing activities occurring below 3,000 ft. are included in emissions estimates. Examples of activities typically occurring below 3,000 ft. include those involving helicopter platforms such as mine warfare, surface warfare, and anti-submarine warfare training and testing activities. The number of all training and testing activities and the estimated time spent above or below 3,000 ft. for calculation purposes is included in the air quality emissions estimates presented in Appendix C (Air Quality Emissions Calculations and Record of Non-Applicability).

The types of aircraft identified include the typical aircraft platforms that conduct a particular training or testing exercise (or the closest surrogate when information is not available), including range support

aircraft (e.g., non-Navy commercial air services). Estimates of future aircraft sorties are based on evolutionary changes in the Navy's force structure and mission assignments. Where there are no major changes in types of aircraft, future activity levels are estimated from the distribution of baseline activities. The types of aircraft used in each training or testing activity along with hours operated in the mission activity, as well as data on landings and take-offs from ships, and numbers of sorties flown by such aircraft are presented in Appendix C (Air Quality Emissions Calculations).

Several testing activities are similar to training activities, and therefore similar assumptions were made for such activities in terms of aircraft type, altitude, and flight duration. Table 2.3-3 lists Naval Air Systems Command testing activities similar to certain training activities. Where aircraft testing activities were dissimilar to training activities, assumptions for time on ranges, and landing and takeoff information were derived by Navy subject matter experts.

Air pollutant emissions from aircraft were primarily estimated based on the training and testing hours provided by subject matter experts, as well as emission indices published in the Navy's Aircraft Environmental Support Office Memorandum Reports for individual aircraft categories. When Aircraft Environmental Support Office emission factor data were not available, emission factors were obtained from other published sources.

The emissions calculations performed for each alternative conservatively assume that each aircraft training and testing activity listed in Tables 2.3-1 to 2.3-4 is separately conducted. In practice, a testing activity may be conducted during a training flight. It is also probable that two or more training activities may be conducted during one flight (e.g., chaff or flare exercises may occur during electronic warfare activities; or air-to-surface gunnery and air-to-surface bombing activities may occur during a single flight operation). Conservative assumptions may produce elevated aircraft emissions calculations but account for the possibility, however remote, that each aircraft training and testing activity is separately conducted.

3.1.1.4.2 Military Vessel Activities

Military vessel traffic in the Study Area includes military ships and smaller boats providing services for military training and testing activities. The methods for estimating military ship emissions involve evaluating the type of activity, generating the average steaming hours for ships in each operational area, both within state waters and beyond state waters. This was done to create annual averages for the years 2010 through 2015. The average annual hours were used for Alternative 1. Alternative 1 reflects a representative year of training to account for the natural fluctuation of training cycles and deployment schedules that generally influence the maximum level of training that may occur year after year in any 5-year period. For Alternative 2, the year with the highest number of operational hours (2011) was selected as the year to represent maximum operations. Alternative 2 reflects the maximum number of training activities that could occur within a given year and assumes that the maximum level of activity would occur every year over any 5-year period. For both alternatives, the hourly data was used with data from the Naval Sea Systems Command database, Navy and Military Sealift Command Marine Engine Fuel Consumption and Emission Calculator to calculate the emissions from the propulsion and onboard generation systems. Data from the calculator included emission factors for each type of propulsion and type of onboard generator by ship type, as well as the fuel used and applicable power levels. The types of ships and numbers of activities for Alternatives 1 and 2 are derived from range records and Navy subject matter experts regarding ship participant data. Estimates of future ship activities are based on anticipated evolutionary changes in the Navy's force structure and mission

assignments. Where there are no major changes in types of ships, estimates of future activities are based on the historical distribution of ship activities. The resulting calculations provided information on the time spent at each power level in each part of the Study Area, emission factors for that power level (in pounds of pollutant per hour), and total emissions for each marine vessel for each operational type and mode.

Small boat emissions were estimated based on activity data provided by the Navy, which included the type and number of small boats, locations, and total number of hours running. Each alternative conservatively assumes that small boat training and testing activities are separately conducted and separately produces emissions. In practice, multiple training/testing activities may be conducted during one training/testing event. Conservative assumptions may produce elevated vessel emissions calculations but account for the possibility, however remote, that each training and testing activity is separately conducted.

Emissions factor data came from the Naval Sea Systems Command database, Navy and Military Sealift Command Marine Engine Fuel Consumption and Emission Calculator. For non-road engines, 100 percent of all of the particulate matter less than or equal to 10 microns in diameter from gasoline and diesel-fueled engines is assumed to be particulate matter less than or equal to 2.5 microns in diameter (U.S. Environmental Protection Agency, 2010).

3.1.1.4.3 Submarine Activities

No U.S. submarines burn fossil fuel under normal operating conditions. However, testing of the emergency diesel generator or training practice of emergency conditions can result in the generator running. For this reason, the emergency generator emissions have been included for submarines, conservatively estimated for steaming hours. This overestimates the actual emissions, but ensures the occasional running of emergency diesel generators is documented as an emission.

3.1.1.4.4 Naval Gunfire, Missiles, Bombs, Other Munitions, and Military Expended Material

Naval gunfire, missiles, bombs, and other types of munitions used in training and testing activities emit air pollutants. To estimate the amounts of air pollutants emitted by munitions during their use, the numbers and types of munitions used during training or testing activities are first totaled. Then generally accepted emissions factors (U.S. Environmental Protection Agency, 2009a) for criteria air pollutants are applied to the total amounts. Finally, the total amounts of air pollutants emitted by each munition type are summed to produce total amounts of each criteria air pollutant under each alternative.

3.1.1.5 Climate Change

Greenhouse gases are compounds that contribute to the greenhouse effect—a natural phenomenon in which gases trap heat within the lowest portion of the earth’s atmosphere (surface-troposphere system), causing heating (radiative forcing) at the surface of the earth. The primary long-lived greenhouse gases directly emitted by human activities are carbon dioxide, methane, nitrous oxide, hydrofluorocarbons, perfluorocarbons, nitrogen trifluoride, and sulfur hexafluoride. Carbon dioxide, methane, and nitrous oxide occur naturally in the atmosphere. These gases influence the global climate by trapping heat in the atmosphere that would otherwise escape to space. The heating effect from these gases is considered the primary cause of the global warming observed over the last 50 years (U.S. Environmental Protection Agency, 2009b). Global warming and climate change affect many aspects of the environment.

The administrator of the USEPA determined that six greenhouse gases in combination endanger both the public health and the public welfare of current and future generations. The USEPA specifically identified carbon dioxide, methane, nitrous oxide, hydrofluorocarbons, perfluorocarbons, and sulfur hexafluoride as greenhouse gases (U.S. Environmental Protection Agency, 2009c).

To estimate global warming potential, which is the heat trapping capacity of a gas, the United States quantifies greenhouse gas emissions using the 100-year timeframe values established in the Intergovernmental Panel on Climate Change Fourth Assessment Report (Intergovernmental Panel on Climate Change, 2007), in accordance with United Nations Framework Convention on Climate Change (United Nations Framework Convention on Climate Change, 2013) reporting procedures. All global warming potentials are expressed relative to a reference gas, carbon dioxide, which is assigned a global warming potential equal to 1. Six other primary greenhouse gases have global warming potentials: 25 for methane, 298 for nitrous oxide, 124 to 14,800 for hydrofluorocarbons, 7,390 to greater than 17,340 for perfluorocarbons, 17,200 for nitrogen trifluoride, and up to 22,800 for sulfur hexafluoride. To estimate the carbon dioxide equivalency of a non-carbon dioxide greenhouse gas, the appropriate global warming potential of that gas is multiplied by the amount of the gas emitted. All seven greenhouse gases are multiplied by their global warming potential and the results are added to calculate the total equivalent emissions of carbon dioxide. The dominant greenhouse gas emitted is carbon dioxide, mostly from fossil fuel combustion (85.4 percent) (U.S. Environmental Protection Agency, 2016c). Weighted by global warming potential, methane is the second largest component of emissions, followed by nitrous oxide. Global warming potential-weighted emissions are presented in terms of equivalent emissions of carbon dioxide, using units of metric tonnes. The Proposed Action is anticipated to release greenhouse gases to the atmosphere. These emissions are quantified (primarily using methods elaborated upon in the Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990 – 2014) for the proposed Navy training and testing in the Study Area, and estimates are presented in Chapter 4 (Cumulative Impacts) (U.S. Environmental Protection Agency, 2016c).

The potential effects of proposed greenhouse gas emissions are by nature global and may result in cumulative impacts because most individual sources of greenhouse gas emissions are not large enough to have any noticeable effect on climate change. Therefore, the impact of proposed greenhouse gas emissions to climate change is discussed in the context of cumulative impacts.

3.1.1.6 Other Compliance Considerations, Requirements, and Practices

Executive Order 13834, *Executive Order Regarding Efficient Federal Operations*, issued on May 17, 2018, establishes policy for federal agencies to prioritize actions that reduce waste, cut costs, enhance the resilience of Federal infrastructure and operations, and enable more effective accomplishment of their missions..

In January 2018, the Department of Defense (DoD) published the results of a global screening level assessment of installation vulnerabilities to climate-related security risks with the goal of identifying serious vulnerabilities and developing necessary adaptation strategies. The survey evaluated risk from flooding, extreme temperatures, wind, drought and wildfire.

In June 2014, DoD released the 2014 Climate Change Adaptation Roadmap to document DoD's efforts to plan for the changes that are occurring or expected to occur as a result of climate change. The Roadmap provides an overview and specific details on how DoD's adaptation will occur and describes ongoing efforts (U.S. Department of Defense, 2014). The Navy is committed to improving energy security and environmental stewardship by reducing reliance on fossil fuels. The Navy is actively developing and

participating in energy, environmental, and climate change initiatives that will increase use of alternative energy and reduce emissions of greenhouse gases. The Navy has adopted energy, environmental, and climate change goals. These goals include increasing alternative energy use Navy-wide to 50 percent by 2020; reducing non-tactical petroleum use; ensuring environmentally sound acquisition practices; ensuring environmentally compliant operations for ships, submarines, aircraft, and facilities operated by the Navy; and implementing applicable elements of the Climate Change Adaptation Roadmap.

Equipment used by military units in the Study Area, including ships and other marine vessels, aircraft, and other equipment, are properly maintained and fueled in accordance with applicable Navy requirements. Operating equipment meets federal and state emission standards, where applicable.

3.1.2 AFFECTED ENVIRONMENT

3.1.2.1 General Background

3.1.2.1.1 Region of Influence


The region of influence for air quality is a function of the type of pollutant, emission rates of the pollutant source, proximity to other emission sources, and local and regional meteorology. Figure 3.1-1 through Figure 3.1-4 present maps of the nonattainment and maintenance areas in the vicinity of the Study Area. For inert pollutants (all pollutants other than ozone and its precursors), the region of influence is generally limited to a few miles downwind from the source. For a photochemical pollutant such as ozone, however, the region of influence may extend much farther downwind. Ozone is a secondary pollutant formed in the atmosphere by photochemical reactions of previously emitted pollutants, or precursors (volatile organic compounds and nitrogen oxides). The maximum impacts of precursors on ozone levels tend to occur several hours after the time of emission during periods of high solar load, and may occur many miles from the source. Ozone and ozone precursors transported from other regions can also combine with local emissions to produce high local ozone concentrations. Therefore, the region of influence for air quality includes the Study Area as well as adjoining land areas several miles inland, which may from time to time be downwind from emission sources associated with the Proposed Action.

3.1.2.2 Sensitive Receptors

Identification of sensitive receptors is part of describing the existing air quality environment. Sensitive receptors are individuals in residential areas, schools, parks, hospitals, or other sites for which there is a reasonable expectation of continuous human exposure during the timeframe coinciding with peak pollution concentrations. On the oceanic portions of the Study Area, crews of commercial vessels and recreational users of the northern Atlantic Ocean and Gulf of Mexico could encounter the air pollutants generated by the Proposed Action. Few such individuals are expected to be present and the duration of substantial exposure to these pollutants is limited because the areas are cleared of nonparticipants before event commencement. These potential receptors are not considered sensitive.


3.1.2.2.1 Climate of the Study Area

The climatic conditions in the Study Area provide background on factors influencing air quality. Climate zones within the Study Area vary with latitude or region. For air quality, the Study Area can be divided into four areas: the North Atlantic Region (Arctic region to Nova Scotia), the Mid-Atlantic Region (Maine to Virginia), the Southeast Atlantic Region (North Carolina to southern Florida) and the Gulf of Mexico Region (southern Florida to Texas).




Notes: AFTT: Atlantic Fleet Training and Testing; OPAREA: Operation Area; VACAPES: Virginia Capes; PM_{2.5}: particulate matter less than or equal to 2.5 microns in diameter; PM₁₀: particulate matter less than or equal to 10 microns in diameter.

Figure 3.1-1: Applicable Nonattainment and Maintenance Areas in USEPA Region 1 and 2




Notes: AFTT: Atlantic Fleet Training and Testing; OPAREA: Operation Area; VACAPES: Virginia Capes; PM_{2.5}: particulate matter less than or equal to 2.5 microns.

Figure 3.1-2: Applicable Nonattainment and Maintenance Areas in USEPA Region 3



Notes: AFTT: Atlantic Fleet Training and Testing; OPAREA: Operation Area; SO₂: sulfur dioxide; Pb: lead.

Figure 3.1-3: Applicable Nonattainment and Maintenance Areas in USEPA Region 4



Notes: AFTT: Atlantic Fleet Training and Testing; OPAREA: Operation Area; SO₂: sulfur dioxide

Figure 3.1-4: Applicable Nonattainment and Maintenance Areas in USEPA Region 6

The climate is arctic near the 65-degree north latitude line and tropical at the 20-degree north latitude line, but most activities and their potential effects would occur in the northern temperate to subtropical climate zones between Maine, Florida, and the Gulf Coast.

The climate of the offshore Atlantic Ocean and adjacent land areas is influenced by the temperatures of the surface waters and water currents as well as by wind blowing across the water. Offshore climates are moderate and seldom have extreme seasonal variations because the ocean is slow to change temperature. Ocean currents of the Atlantic Ocean (i.e., Labrador, Gulf Stream, North Atlantic Drift, Canary, and North Equatorial) influence climate by moving warm and cold water between regions. Adjacent land areas are affected by wind that is cooled or warmed when blowing over these currents. In addition to its influence on temperature, the wind moves evaporated moisture from the ocean to adjacent land areas and is a major source of rainfall.

With the advent of human induced climate change, spatial and temporal variations in weather patterns have emerged or have become more pronounced. Very heavy precipitation events have increased across the eastern half of the United States, with the most pronounced increase involving the mid-Atlantic and New England states (Melillo et al., 2014). Other changes apparent along the eastern seaboard include the rising incidence of heat waves and their extended duration and coastal flooding due to sea level rise and storm surge. In the South and along the Gulf Coast, the incidence of extreme storms, such as hurricanes, continues to rise. These changes to weather patterns have long-term consequences for regional climates and the flora and fauna of the regions.

3.1.2.2.1.1 Newfoundland-Labrador Shelf and Scotian Shelf

The Newfoundland-Labrador Shelf and Scotian Shelf are not connected to the continental United States and do not include state waters, but do fall within the AFTT Study Area. This area does not fall under the purview of the Clean Air Act and, therefore, is not included in the regulatory air quality analysis. In the North Atlantic (Newfoundland-Labrador Shelf and Scotian Shelf) winter begins (when daily temperatures average 32° Fahrenheit [° F]) as early as mid-August in the Labrador Sea or as late as October 1 off the coast of the island of Newfoundland (Canadian Coast Guard, 2010). Winter ends in this region in mid-June. Sea ice begins to grow shortly after the onset of winter as average sea temperatures reach 29° to 35° F. Polar lows usually occur during the fall, winter, and early spring.

3.1.2.2.1.2 Mid-Atlantic United States Continental Shelf

Along the coasts of Maine to Virginia, the most frequent wind directions measured by buoys are from the west or west-northwest, but wind can come from any direction (National Oceanic and Atmospheric Administration, 2017). The average wind speeds are between 12.4 and 16.2 miles per hour (mph). Wind speeds are typically lowest in July at 9.0 to 12.1 mph, and highest in January at 15.7 to 20.0 mph.

Annual average air temperature ranges from 47° to 68° F along the coast of Maine to Virginia (National Oceanic and Atmospheric Administration, 2017). Seasonal variations in temperature are greatest during the winter months. In January and February, the ambient temperature average ranges from 28° F along the coast of Maine to 52° F in Virginia. During the warmer months, there is little daily variation in temperature. In August, the average temperature is 75° F along the coast of this region.

Along the coasts of Maine to Virginia, precipitation is frequent and abundant but occurs evenly throughout the year (Minerals Management Service, 2007a). Average annual rainfall along the Atlantic coast is about 42 inches (in.). Rainfall in the warmer months is usually associated with cloud systems that produce showers and thunderstorms. Winter rains are associated with the passage of frontal

systems through the eastern seaboard. Precipitation also falls as snow along the coasts of Maine to the highlands of Virginia. The highest snowfall among coastal U.S. areas within the Study Area occurs in Portland, Maine, with a maximum yearly average of 62.4 in.

3.1.2.2.1.3 Southeast United States Continental Shelf

Off the coast of North Carolina, the prevailing winds are from south to southwest, with average wind speeds between 13 to 16 mph. Off the coasts of South Carolina and Georgia, the prevailing wind direction is from south to southwest, and from southeast to east-southeast off of Florida. Average wind speeds range from 12 to 14 mph and wind speeds exhibit smaller monthly variations than northern coastal states.

Annual average air temperatures range from 70° to 75° F along the coast of the Southeast U.S. Continental Shelf (National Oceanic and Atmospheric Administration, 2017). In January and February, ambient temperatures average 55°F along the coast of the Southeast U.S. Continental Shelf. During the warmer months, there is little daily variation in temperature. In August, average temperatures are 83° F along the coast of this region. Air temperatures over the southern coast and offshore Atlantic Ocean have smaller daily and seasonal ranges than temperatures over inland areas because the ocean, which is slow to change temperature, has a stabilizing influence on ocean and coastal atmospheric temperatures.

At various locations along the Atlantic coast, fog occurs occasionally in the cooler months as a result of warm, moist air from the Gulf of Mexico blowing over cool land or water surfaces. The poorest visibility occurs from November through April. During periods of air stagnation, industrial pollution and agricultural burning also can affect visibility.

In the Southeast U.S. Continental Shelf coastal areas (generally from North Carolina to Florida), precipitation is frequent and abundant throughout the year, but tends to peak in the summer months.

Hurricanes develop in the southern part of the Atlantic Ocean. Hurricane season in the Atlantic Ocean runs from June to November, with a peak in mid-September. Most storms form in warm waters several hundred miles north of the equator. Once a tropical system forms, it usually travels west and slightly north while strengthening. Many storms curve to the northeast near the Florida peninsula. The Atlantic basin averages about 10 storms of tropical storm strength or greater per year; about half reach hurricane level (National Oceanic and Atmospheric Administration, 2005). Storms weaken as they encounter cooler water, land, or vertical wind shear, sometimes slowing to an extra-tropical storm, mostly affecting northern Atlantic coastal areas.

3.1.2.2.1.4 Gulf of Mexico

The climate of the Gulf of Mexico is influenced mainly by the clockwise circulation around the semi-permanent area of high barometric pressure commonly known as the Bermuda High (Minerals Management Service, 2002). The Gulf of Mexico is southwest of this center of circulation. This high-pressure system results in a predominantly southeasterly wind flow in the Gulf of Mexico. Two important classes of storms occasionally occur with this circulation pattern. During the winter months, cold fronts associated with cold air masses from land influence the northern coast of the Gulf of Mexico. Behind the fronts, strong north winds bring drier air into the region. Secondly, hurricanes may develop in or migrate into the Gulf of Mexico during the warmer months. These storms may affect any area of the Gulf of Mexico and substantially change the local wind circulation around them. In coastal areas, the sea breeze may become the primary circulation feature during the summer months. Conversely, land breezes (particularly at night) transport air pollutants from land to offshore areas. Locally, the land

breeze diminishes as more heat is retained within large, growing coastal cities (National Science Foundation, 2011). In general, however, the subtropical maritime climate is the dominant feature driving all aspects of the weather in this region. As a result, the climate shows very little daily or seasonal variation (Minerals Management Service, 2002).

Average air temperatures at Gulf of Mexico coastal locations (Texas to Florida) vary with latitude and exposure. Air temperatures range from highs in the summer of 88° to 96° F to lows in the winter of 37° to 59° F (Minerals Management Service, 2002). Temperatures depend on the frequency and intensity of polar air masses from the north. Air temperatures over the open waters of the Gulf of Mexico are more moderate and have smaller daily and seasonal temperature ranges than land temperatures because the Gulf of Mexico is slow to change temperature (Minerals Management Service, 2007b). The average temperature over the center of the Gulf of Mexico is about 84° F in the summer and between 63° to 73° F in the winter (Minerals Management Service, 2007b).

In the Gulf of Mexico portion of the Study Area, precipitation is frequent and abundant throughout the year (Minerals Management Service, 2002). Stations along the entire Gulf Coast record the highest precipitation values during the warmer months of the year. The warmer months usually have cloud systems that produce showers and thunderstorms; however, these thunderstorms rarely cause any damage or have hail (Minerals Management Service, 2002). The month of maximum rainfall for most locations in the Gulf of Mexico is July. Winter rains often come with frontal systems passing through the area. Rainfall is generally light, steady, and relatively continuous, often lasting several days. Snowfall is rare, and when snow or sleet does occur, it usually melts on contact with the ground. The chance for snow or sleet decreases with distance from shore, rapidly reaching zero.

Hurricanes affecting the Gulf of Mexico form near the equator in the Atlantic Ocean, the Caribbean Sea, and the Gulf of Mexico (Minerals Management Service, 2002). Data from 1886 to 1986 show that almost half (44.5 percent) of these hurricanes, or 3.7 storms per year, will affect the Gulf of Mexico (Minerals Management Service, 2002).

3.1.2.3 Existing Air Quality

As a whole, the air quality of the Study Area is very good. As shown in Figure 3.1-1 through Figure 3.1-3, most nonattainment and maintenance areas in the eastern half of the continental United States are in the northeastern states. Many are located in inland, urban, industrialized areas. This limited geographical extent with regard to potential air pollution results from the relatively low number of air pollutant sources, size, and topography of the Study Area, and prevailing meteorological conditions. In general, the coastal counties of the lower-middle and southern Atlantic as well as the Gulf of Mexico, including the Hampton Roads Intrastate area (in the vicinity of Naval Station Norfolk on Figure 3.1-2), are in attainment of the National Ambient Air Quality Standards. Being in attainment means that the areas maintain air quality better than the National Ambient Air Quality Standards.

Some other coastal areas, however, are either in nonattainment or are a designated maintenance area for one or more of the criteria pollutants. These designations are based on air quality data collected from monitors at locations in urban and rural setting, as well as modeling. Based on available information, the USEPA designates an area as attainment, maintenance, nonattainment, or if there is a lack of available monitoring data for the area, it may be designated unclassifiable. Nonattainment and maintenance designations range from as small as a single location to large multi-state regions. Table 3.1-3 identifies the nonattainment and maintenance areas that are adjacent to the Study Area.

Table 3.1-3: Nonattainment and Maintenance Areas Adjacent to Study Area

<i>Area Name</i>	<i>Coastal Locations Included</i>	<i>Designation</i>
USEPA Regions 1 & 2		
Central New Hampshire, NH	Rockingham County (p), Hillsborough County (p)	2010 SO ₂ (n)
Greater Connecticut	New London County	Ozone (n-moderate)
Hartford –New Britain- Middletown, CT	Middlesex County CT (p)	CO (m)
New Haven-Meriden-Waterbury, CT	New Haven County CT	CO (m)
New York-Northern New Jersey- Long Island, NY-NJ-CT	Fairfield, New Haven & Middlesex Counties (CT); Bronx, Kings, Nassau, New York, Queens, Richmond, Rockland, Suffolk, & Westchester Counties (NY); Bergen, Essex, Hudson, Union, Middlesex & Monmouth Counties (NJ)	Ozone (n-moderate)
	Fairfield & New Haven Counties (CT); Bronx, Kings, Nassau, New York, Queens, Richmond, Rockland, Suffolk, & Westchester Counties (NY); Bergen, Essex, Hudson, Union, Middlesex & Monmouth Counties (NJ)	1997 PM _{2.5} (m) and 2006 PM _{2.5} (m)
	New Haven County CT (p)	PM ₁₀ (m)
	New York County NY	PM ₁₀ (n)
	Fairfield County CT (p)	CO (m)
Philadelphia-Wilmington-Atlantic City, PA-NJ-MD-DE	Atlantic, Cape May & Ocean Counties	Ozone (n-marginal)
USEPA Region 3		
Seaford, DE	Sussex County	Ozone (n-marginal)
USEPA Region 4		
Nassau County, FL	Nassau County, FL (p)	2010 SO ₂ (n)
Hillsborough County, FL	Hillsborough County, FL (p)	2010 SO ₂ (n)
	Tampa, FL (p)	2008 Lead
USEPA Region 6		
Saint Bernard Par LA	Saint Bernard Parish, LA	2010 SO ₂ (n)
Houston-Galveston-Brazoria, TX	Brazoria, Chambers, Galveston Counties, TX	Ozone (n-moderate)

Source: (U.S. Environmental Protection Agency, 2017b)

Notes: (p) means partial; (n) means nonattainment; (m) means maintenance

CO: carbon monoxide; PM₁₀: particulate matter ≤ 10 microns in diameter; PM_{2.5}: particulate matter ≤ 2.5 microns in diameter; SO₂: sulfur dioxide

The Greater Connecticut area is designated as moderate nonattainment for ozone. Table 3.1-4 lists Study Area pierside locations and the attainment status for each.

Table 3.1-4: Pierside and Coastal Activity Locations and Their Area’s Attainment Status

<i>Pierside Location</i>	<i>Designated Area</i>	<i>National Ambient Air Quality Standards Attainment Status</i>
Portsmouth Naval Shipyard, Kittery Maine; Shipyard – Bath, Maine	Metropolitan Portland/ Cumberland County	Attainment of all applicable standards
Naval Undersea Warfare Center, Division, Newport, Newport, Rhode Island	Providence (all of RI), RI	Attainment of all applicable standards
Naval Submarine Base New London; Groton, Connecticut Shipyard – Groton, Connecticut and Thames River	Greater Connecticut, CT	Moderate nonattainment of the 8-hour ozone standard Attainment of all other applicable standards
Naval Station Norfolk, Norfolk, Virginia; Joint Expeditionary Base Little Creek-Fort Story, Virginia Beach, Virginia; Norfolk Naval Shipyard, Portsmouth, Virginia; Shipyard – Newport News, Virginia; Broad Bay; York River; James River and Tributaries	Hampton Roads Intrastate	Attainment of all applicable standards
Cooper River; Charleston Pier, South Carolina	Charleston County	Attainment of all applicable standards
Naval Submarine Base Kings Bay, Georgia	Camden County	Attainment of all applicable standards
Naval Station Mayport, Jacksonville, Florida; St. Johns River, Florida	Duval County	Attainment of all applicable standards
Port Canaveral, Cape Canaveral, Florida	Brevard County	Attainment of all applicable standards
Saint Andrew Bay, Florida	Bay County	Attainment of all applicable standards
Shipyard – Pascagoula, Mississippi	Jackson County	Attainment of all applicable standards

Source: 40 CFR part 81, Subpart C and Green Book Nonattainment and Maintenance Areas (U.S. Environmental Protection Agency, 2017b)

Figure 3.1-1 through Figure 3.1-4 show the nonattainment and maintenance areas that are within or adjacent to the AFTT Study Area.

3.1.2.3.1 Other Air Basins Adjacent to the Study Area

A substantial portion (over 70 percent) of all AFTT EIS/OEIS training and testing activities occur within the range complexes, which are adjacent to coastal attainment areas but located beyond state waters. The remaining 30 percent are largely conducted well offshore and a small percentage is performed in areas offshore of coastal nonattainment or maintenance areas. These areas include stretches of coastal areas of the northeast, areas adjacent to Nassau County, Florida, the Tampa area, the New Orleans area, and coastal areas around Houston. The migration of emissions from offshore sources to land is well-documented. In 1997, the International Maritime Organization adopted Annex VI, Regulations for the Prevention of Air Pollution from ships. These regulations were instituted for the commercial maritime industry due to recognition of the impact of vessel emissions, which can travel hundreds of miles, on coastal receptors and further inland. These emissions are particularly significant around the large ports

on the coast of the United States, which include New York/New Jersey, Philadelphia, Baltimore, Norfolk, Charleston, Savannah, Jacksonville, Miami, South Louisiana, and Houston (U.S. Maritime Administration, 2016).

In addition to the Operational Areas (OPAREAs) and other areas further out to sea, there are also activities that occur within state waters. Vessels traverse state water during ingress/egress to OPAREAs and other Study Area locations further offshore. There are also training activities in particular that occur in coastal areas, including riverine and bay locations. The area of greatest activity is in the lower Chesapeake Bay and in tributaries to the bay, primarily the James and York Rivers in Virginia. Activities in Narragansett Bay are associated with the Naval Undersea Warfare Center Division, Newport in Rhode Island. Additional areas where training or testing occurs within state waters include the St. Johns River near Naval Station Mayport, Florida, Port Canaveral, Florida, St. Andrew Bay near Naval Support Activity Panama City, Florida and the Cooper River near Charleston, South Carolina. Of these, only Naval Station Mayport is located in an Air Quality Control Region with a nonattainment designation within its borders.

3.1.3 ENVIRONMENTAL CONSEQUENCES

This section evaluates how and to what degree the activities described in Chapter 2 (Description of Proposed Action and Alternatives) potentially impact air quality within the Study Area. The air quality stressors vary in intensity, frequency, duration, and location within the Study Area. The stressor applicable to air quality in the Study Area is analyzed below:

- **Criteria Air Pollutants**

In this analysis, criteria air pollutant emissions estimates were calculated for vessels, aircraft, and munitions. For each alternative, emissions estimates were developed by range complex and other training or testing locations and totaled for the Study Area. Additionally, state waters emissions are separately analyzed for air quality impacts. Details of the emission estimates are provided in Appendix C (Air Quality Emissions Calculations and Record of Non-Applicability). Hazardous air pollutants are analyzed qualitatively in relation to the prevalence of the sources emitting hazardous air pollutants during training and testing activities.

Estimating Emissions of Criteria Air Pollutants

The potential impacts of criteria air pollutants are evaluated by first estimating the emissions from training and testing activities in the Study Area for each alternative. These estimates are then used to determine the potential impact of the emissions on the attainment status of the adjacent designated air quality area.

The estimate of criteria air pollutant emissions for each alternative is categorized by region (e.g., by range complex or testing range) so that differences in background air quality, atmospheric circulation patterns, regulatory requirements, and sensitive receptors can be addressed. An overall estimate of air pollutant emissions for Navy training and testing activities in the Study Area under each alternative is also provided. Under Alternative 1, emissions were based on the average number of training and testing activities anticipated, based on the prior 6 years of data. Under Alternative 2, emissions were based on the anticipated maximum number of training and testing activities. For vessel operations, the maximum was based on the operations that occurred in 2011 the year of the highest number of operations in the range 2010 – 2015. While this represented the year of most total operations, the number of operations involving specific vessels in the individual operational areas may or may not have been higher than the

average number used in Alternative 1. These individual variances do not change the overall result of greater total operations when accounting for all vessels in all regions under Alternative 2.

Table 3.1-5 and Table 3.1-7 present the total emissions for the proposed training and testing activity locations under each alternative. Table 3.1-6 and Table 3.1-8 present the emissions specific to the Jacksonville nonattainment/maintenance area with a comparison to the General Conformity *de minimis* thresholds to assess the applicability of the General Conformity Regulations to the proposed action in this area.

The analysis below is organized by Alternative and provides both the NEPA impact analysis, and where appropriate, a separate discussion related to General Conformity requirements.

3.1.3.1 Impacts from Air Emissions Under Alternative 1

3.1.3.1.1 NEPA Impacts from Air Emissions under Alternative 1

Table 3.1-5 presents the total estimated emission results under Alternative 1 for each operational region in the Study Area and includes all emissions generated, regardless of proximity to the coastline. Most of these emissions occur beyond state waters. For Virginia Capes Range Complex, the use of vessels within the state waters is up to 2 percent, and in the Jacksonville Range Complex, the use of vessels within state waters is up to 1 percent.

The subsections that follow evaluate the emissions, both within the entire Study Area, and in state waters within the Study Area, under NEPA.

Table 3.1-5: Estimated Annual Air Pollutant Emissions from Activities Occurring within the AFTT Study Area, Alternative 1

Range Complex	Emissions by Air Pollutant (TPY)					
	VOC	CO	NO _x	SO _x	PM ₁₀	PM _{2.5}
Northeast	6.0	25.0	143.4	28.3	9.7	9.7
Virginia Capes	121.9	939.8	3,722.7	984.1	190.7	190.7
Cherry Point	43.8	348.8	918.7	172.7	42.3	42.3
Jacksonville	49.4	485.7	1,069.8	306.6	71.5	71.5
Key West	2.8	12.2	76.7	12.9	5.4	5.4
Gulf of Mexico	8.3	122.3	416.5	108.6	25.2	25.2
Outside Range Complex Areas	53.7	332.8	1,683.1	383.5	55.5	55.5

Notes: CO: carbon monoxide; NO_x: oxides of nitrogen; VOC: volatile organic compounds; SO_x: sulfur oxides; PM₁₀: particulate matter less than or equal to 10 microns in diameter; PM_{2.5}: particulate matter less than or equal to 2.5 microns in aerodynamic diameter; TPY: tons per year.

A significant portion of the Study Area activities would occur well offshore. While pollutants emitted in the Study Area under Alternative 1 may at times be carried ashore by winds, most training and testing activities would occur more than 12 NM offshore, and natural mixing would substantially disperse pollutants before they reach the coastal land mass. The contributions of air pollutants generated in the Study Area to the air quality in onshore areas are unlikely to measurably add to existing onshore pollutant concentrations because of the distances these offshore pollutants would be transported and their substantial dispersion during transport.

3.1.3.1.2 General Conformity Analysis under Alternative 1 in Northeast Areas Designated Nonattainment or Maintenance

In the northeast, the primary areas where air pollution has resulted in designation of nonattainment or maintenance areas lie in the New York-Northern New Jersey-Long Island, NY-NJ-CT Air Quality Control Region (U.S. Environmental Protection Agency, 1972) (see Figure 3.1-1) which is moderate nonattainment for ozone, a maintenance area for particulate matter less than or equal to 2.5 microns in diameter, and includes a maintenance area for particulate matter less than or equal to 10 microns in diameter. A portion of the Eastern Connecticut Intrastate Control Region is also designated as moderate nonattainment for ozone. A very small area of coastal New Hampshire is nonattainment for sulfur dioxide, and there is a small area of ozone nonattainment in the coastal counties of New Jersey as well as near the coast at Seaford, Delaware. Activities in state waters are not scheduled to occur in any of these nonattainment or maintenance areas. The primary location where activities in state waters occur is at Naval Undersea Warfare Center Newport and Narragansett Bay, both of which are in Rhode Island, an area in attainment for all pollutants.

3.1.3.1.3 General Conformity Analysis under Alternative 1 in Jacksonville Florida Areas Designated Nonattainment or Maintenance

In the southeast, the area where air pollution has resulted in designation of a coastal nonattainment or maintenance area lies in Nassau County, Florida, which is just north of Jacksonville (see Figure 3.1-3). Both of these counties are in the Jacksonville (Florida)-Brunswick (Georgia) Interstate Air Quality Control Region. A portion of Nassau County is nonattainment for sulfur dioxide which will require that a General Conformity applicability analysis be performed to determine if a formal General Conformity Determination is required. Table 3.1-6 presents the estimated state waters emissions and their relevance to applicable General Conformity thresholds.

Table 3.1-6: Estimated Annual Air Pollutant Emissions from Activities Occurring in State Waters in the Jacksonville, Florida Area, Alternative 1

	<i>Emissions by Air Pollutant (TPY)</i>					
	<i>VOC</i>	<i>CO</i>	<i>NO_x</i>	<i>SO_x</i>	<i>PM₁₀</i>	<i>PM_{2.5}</i>
Nassau, FL SO ₂ Nonattainment Area						
Total Emissions from all Sources	1.3	6.6	24.6	4.9	1.3	1.3
General Conformity Thresholds	NA	NA	NA	100	NA	NA
Exceedance?	NA	NA	NA	No	NA	NA

Notes: Table includes criteria pollutant precursors (e.g., VOC). Individual values may not add exactly to total values due to rounding.

CO: carbon monoxide; NO_x: nitrogen oxides; PM_{2.5}: particulate matter less than or equal to 2.5 microns in diameter;

PM₁₀: particulate matter less than or equal to 10 microns in diameter; SO_x: sulfur oxides; TPY: tons per year; VOC: volatile organic compounds

Sulfur dioxide emissions in state waters associated with AFTT activities would be below the General Conformity *de minimis* thresholds. As a result, no further analysis of conformity is required and a Record of Non-Applicability, located in Appendix C, was prepared in accordance with Navy guidance.

3.1.3.1.4 General Conformity Analysis under Alternative 1 in the Gulf of Mexico Areas Designated Nonattainment or Maintenance

In the Gulf of Mexico, the primary areas where air pollution has resulted in designation of nonattainment or maintenance areas lie in Hillsborough County, Florida (see Figure 3.1-1) which is

nonattainment for sulfur dioxide and lead; Saint Bernard Parish, Louisiana, which is also nonattainment for sulfur dioxide; and the Houston-Galveston-Brazoria ozone nonattainment area. Activities in state waters are not scheduled to occur in any of these nonattainment or maintenance areas. The primary location where state water activities in this region do occur is at Naval Surface Warfare Center Panama City Division, Florida which is in attainment for all pollutants.

3.1.3.1.5 Summary of Impacts from Criteria Pollutants under Alternative 1

While both criteria and hazardous air pollutants emitted in the Study Area under Alternative 1 may at times be carried ashore by prevailing winds, most training and testing activities would occur beyond state water boundaries and natural mixing would substantially disperse pollutants before they reach the boundaries of the adjacent air quality control regions. Additionally, the primary wind pattern moves from shore to offshore (National Aeronautics and Space Administration, 2018). The contributions of air pollutants generated in the Study Area to the air quality in the air quality control regions are unlikely to measurably add to existing onshore pollutant concentrations because of the distances these offshore pollutants would be transported, their substantial dispersion during transport and the intermittent short-term nature of the activities. Therefore, no significant impacts on air quality as a result of criteria pollutants over state waters would occur; and no significant harm to air quality as a result of criteria pollutant emissions beyond state waters would occur. Total direct and indirect emissions of criteria pollutants under Alternative 1 also fall below *de minimis* levels, thus a General Conformity Analysis is not required.

3.1.3.2 Impacts from Air Emissions under Alternative 2

3.1.3.2.1 NEPA Impacts from Air Emissions under Alternative 2

Table 3.1-7 presents the total estimated emission results under Alternative 2 for each operational region in the Study Area and includes all emissions generated, regardless of proximity to the coastline. Most of these emissions occur beyond state waters. For Virginia Capes Range Complex, the use of vessels within the state waters is greater than in other portions of the Study Area.

The subsections that follow evaluate the state waters emissions within the regional areas that include nonattainment or maintenance areas. These emissions are compared to the General Conformity *de minimis* thresholds, and are not specific to specific localities. This conservative approach, then, evaluates all nearshore emissions as potentially occurring in any of the applicable nonattainment or maintenance areas.

Table 3.1-7: Estimated Annual Air Pollutant Emissions from Activities Occurring within the AFTT Study Area, Alternative 2

	<i>Emissions by Air Pollutant (TPY)</i>					
	<i>VOC</i>	<i>CO</i>	<i>NO_x</i>	<i>SO_x</i>	<i>PM₁₀</i>	<i>PM_{2.5}</i>
Northeast	5.5	26.3	120.8	20.3	12.1	12.1
Virginia Capes	118.0	935.9	3,993.9	1,070.8	211.6	211.6
Cherry Point	33.1	185.9	821.5	194.7	39.9	39.9
Jacksonville	61.2	602.8	1,994.2	540.3	88.9	88.9
Key West	0.9	14.3	30.3	10.6	3.7	3.7
Gulf of Mexico	1.7	27.4	59.7	19.7	9.7	9.7
Outside Range Complex Areas	162.4	569.7	4,160.7	656.8	90.2	90.2

Notes: CO: carbon monoxide; NO_x: oxides of nitrogen; VOC: volatile organic compounds; SO_x: sulfur oxides; PM₁₀: particulate matter less than or equal to 10 microns in diameter; PM_{2.5}: particulate matter less than or equal to 2.5 microns in diameter; TPY: tons per year.

A significant portion of the Study Area activities would occur well offshore. While pollutants emitted in the Study Area under Alternative 2 may at times be carried ashore by winds, most training and testing activities would occur more than 12 NM offshore, and natural mixing would substantially disperse pollutants before they reach the coastal land mass. The contributions of air pollutants generated in the Study Area to the air quality in onshore areas are unlikely to measurably add to existing onshore pollutant concentrations because of the distances these offshore pollutants would be transported and their substantial dispersion during transport.

3.1.3.2.2 General Conformity Analysis under Alternative 2 in Northeast Areas Designated Nonattainment or Maintenance

In the northeast, the primary areas where air pollution has resulted in designation of nonattainment or maintenance areas lies in the New York-Northern New Jersey-Long Island, NY-NJ-CT Air Quality Control Region (U.S. Environmental Protection Agency, 1972) (see Figure 3.1-1) which is moderate nonattainment for ozone, a maintenance area for particulate matter less than or equal to 2.5 microns in diameter, and includes a maintenance area for particulate matter less than or equal to 10 microns in diameter. A portion of the Eastern Connecticut Intrastate Control Region is also designated as moderate nonattainment for ozone. A very small area of coastal New Hampshire is nonattainment for sulfur dioxide, and there is a small area of ozone nonattainment near the coast at Seaford, Delaware. State waters activities are not scheduled to occur in any of these nonattainment or maintenance areas. The primary location where state waters activities in this region do occur is at Naval Undersea Warfare Center Division Newport and Narragansett Bay, both of which are in Rhode Island, an area in attainment for all pollutants.

3.1.3.2.3 General Conformity Analysis under Alternative 2 in Jacksonville Florida Areas Designated Nonattainment or Maintenance

In the southeast, the area where air pollution has resulted in designation of a coastal nonattainment or maintenance area lies in the Nassau County, Florida, which is just north of Jacksonville (see Figure 3.1-3). Both of these counties are in the Jacksonville (Florida)-Brunswick (Georgia) Interstate Air Quality Control Region. A portion of this county is nonattainment for sulfur dioxide. Table 3.1-8 presents the estimated nearshore emissions and their relevance to applicable General Conformity thresholds.

Table 3.1-8: Estimated Annual Air Pollutant Emissions from Activities Occurring within 3 NM of shore in the Jacksonville, Florida Area, Alternative 2

	Emissions by Air Pollutant (TPY)					
	VOC	CO	NO _x	SO _x	PM ₁₀	PM _{2.5}
Nassau, FL SO₂ Nonattainment Area						
Total Emissions from all Sources	1.5	9.6	30.7	6.1	1.6	1.6
General Conformity Thresholds	NA	NA	NA	100	NA	NA
Exceedance?	NA	NA	NA	No	NA	NA

Notes: Table includes criteria pollutant precursors (e.g., VOC). Individual values may not add exactly to total values due to rounding.

CO: carbon monoxide; NO_x: nitrogen oxides; PM_{2.5}: particulate matter less than or equal to 2.5 microns in diameter; PM₁₀: particulate matter less than or equal to 10 microns in diameter; SO_x: sulfur oxides; TPY: tons per year; VOC: volatile organic compounds

Sulfur dioxide emissions in state waters that are associated with AFTT activities would be below the General Conformity *de minimis* thresholds. As a result, no further analysis of conformity is required and a Record of Non-Applicability, located in Appendix C, was prepared in accordance with Navy guidance.

3.1.3.2.4 General Conformity Analysis under Alternative 2 in the Gulf of Mexico Adjacent Areas Designated Nonattainment or Maintenance

In the Gulf of Mexico, the primary areas where air pollution has resulted in designation of nonattainment or maintenance areas lie in Hillsborough County, Florida (see Figure 3.1-1) which is nonattainment for sulfur dioxide and lead; Saint Bernard Parish, Louisiana, which is also nonattainment for sulfur dioxide; and the Houston-Galveston-Brazoria ozone nonattainment area. State waters activities are not scheduled to occur in any of these nonattainment or maintenance areas. The primary location where state waters activities in this region do occur is at Naval Surface Warfare Center Panama City Division, Florida which is in attainment for all pollutants. As a result, the General Conformity Regulations do not apply.

3.1.3.2.5 Summary of Impacts from Criteria Pollutants under Alternative 2

While pollutants emitted in the Study Area under Alternative 2 may at times be carried ashore by prevailing winds, most training and testing activities would occur more than 3 NM offshore, and natural mixing would substantially disperse pollutants before they reach the boundaries of the adjacent air quality control regions. The contributions of air pollutants generated in the Study Area to the air quality in the air quality control regions are unlikely to measurably add to existing onshore pollutant concentrations because of the distances these offshore pollutants would be transported and their substantial dispersion during transport. Total direct and indirect emissions of criteria pollutants under Alternative 1 also fall below *de minimis* levels, thus a General Conformity Analysis is not required.

3.1.3.3 Impacts from Air Pollutants under the No Action Alternative

Under the No Action Alternative, training and testing activities associated with the Proposed Action would not be conducted within the AFTT Study Area. Discontinuing training and testing activities in the Study Area under the No Action Alternative would not measurably improve air quality in the Study Area because of the discontinuous nature of the events that constitute the Proposed Action and the fact that most of the air emissions that are generated occur at sea over a wide geographic area. The elimination of the air emissions associated with training activities in the lower Chesapeake Bay and its tributaries may be beneficial to local air quality in this region because it is the area of highest activity in state waters. It should be noted that the air quality in this area already surpasses the National Ambient Air Quality Standards.

3.1.3.4 Greenhouse Gases and Climate Change

Activities conducted as part of the Proposed Action would involve mobile sources using fossil fuel combustion as a source of power. Additionally, the expenditure of munitions could generate greenhouse gas emissions. While the emissions generated by testing and training activities alone would not be enough to cause global warming, in combination with past and future emissions from all other sources they would contribute incrementally to the global warming that produces the adverse effects of climate change.

Greenhouse gas emissions for all of the training and testing activities occurring annually throughout the entire Study Area were calculated using emissions factors provided by the U.S. Navy for aircraft and vessels, and published by the USEPA for munitions. The analysis of greenhouse gas emissions associated with aircraft is limited to those emissions below 3,000 ft. because there is insufficient historical data to document the entire flight path or flight duration of any given aircraft for a specific training or testing event. This is also true for the baseline data so that the totals for the baseline, Alternative 1 and Alternative 2 are comparable. A comparative analysis was performed using the greenhouse gas emission estimates prepared for the Preferred Alternative in the 2013 AFTT Final EIS/OEIS. A net decrease in greenhouse gas emissions would be anticipated compared to the 2013 estimates, with the largest decrease associated with Alternative 1, as indicated in Table 3.1-9.

Table 3.1-9: Total Annual Greenhouse Gas Emissions from All Study Area Training and Testing Activities in Metric Tons per Year

<i>2013 Emission Estimates</i>	<i>Alternative 1 Emissions</i>	<i>Net Change from 2013 Estimates</i>	<i>Alternative 2 Emissions</i>	<i>Net Change from 2013 Estimates</i>
1,360,794	1,027,261	-333,533	1,235,075	-125,719

3.1.4 SUMMARY OF POTENTIAL IMPACTS ON AIR QUALITY

In this analysis, criteria air pollutant and greenhouse gas emissions estimates were calculated for vessels, aircraft, and munitions. For each alternative, emissions estimates were developed by range complex and other training or testing locations and totaled for the Study Area. Details of the emission estimates are provided in Appendix C (Air Quality Emissions Calculations and Example Record of Non-Applicability).

3.1.4.1 Combined Impacts of All Stressors under Alternative 1

Emissions associated with Study Area training and testing activities under Alternative 1 primarily occur beyond the boundary for state waters. For fixed-wing aircraft activities, emissions typically occur above the 3,000 ft. mixing layer. Given these characteristics, the impact on air quality from the combination of these resource stressors are expected to be similar to the impacts on air quality for any of these stressors taken individually without any additive synergistic, or antagonistic interaction. To provide a general comparative analysis, the emissions data for each alternative were compared to the emission data from the Preferred Alternative selected in the 2013 AFTT Final EIS/OEIS. A comparison of estimated emissions under Alternative 1 to the 2013 Preferred Alternative (which was subsequently adopted) indicates that some pollutant emissions would be reduced and others would increase. Emissions of volatile organic compounds, carbon monoxide, and greenhouse gases would decrease. Nitrogen oxides, sulfur dioxide and particulate matter would increase. A significant cause of the increase in nitrogen oxide, particulate matter, and sulfur dioxide emissions is due to the inclusion of riverine and bay testing and training activities, particularly in the Virginia environs. These activities were not analyzed as part of the air quality analysis in the 2013 document.

3.1.4.2 Combined Impacts of All Stressors under Alternative 2

Emissions associated with Study Area training and testing activities under Alternative 2 primarily occur beyond the boundary for state waters. For fixed-wing aircraft activities, emissions typically occur above the 3,000 ft. mixing layer. Given these characteristics, the impact on air quality from the combination of these resource stressors are expected to be similar to the impacts on air quality for any of these stressors taken individually without any additive synergistic, or antagonistic interaction. A comparison of estimated emissions under Alternative 2 in comparison to the baseline indicates that some pollutants emissions would be reduced and others would increase. Emissions of volatile organic compounds, carbon monoxide, and greenhouse gases would decrease. Nitrogen oxides, particulate matter, and sulfur dioxide would increase. A significant cause of the increase in nitrogen oxide and sulfur dioxide emissions is due to the inclusion riverine and bay testing and training activities, particularly in the Virginia environs. These activities were not accounted for in the analyses presented in the phase 2013 AFTT Final EIS/OEIS Proposed Action.

3.1.4.3 Combined Impacts of All Stressors under the No Action Alternative

Training and testing activities associated with the Proposed Action would not be conducted within the AFTT Study Area.

Discontinuing training and testing activities in the Study Area under the No Action Alternative would not measurably improve air quality in the Study Area because of the discontinuous nature of the events that constitute the Proposed Action and the fact that most of the air emissions that are generated occur at sea over a wide geographic area. The elimination of the air emissions associated with training activities in the lower Chesapeake Bay and its tributaries may be beneficial to local air quality in this region because it is the area of highest activity in state waters. It should be noted that the air quality in this area is already better than the National Ambient Air Quality Standards.

References

- Arfsten, D. P., C. L. Wilson, and B. J. Spargo. (2001). Human environmental health issues related to use of radio frequency chaff. *Navy Medicine*, 92(5), 12–16.
- Canadian Coast Guard. (2010). *Chapter 3: Ice and Weather Environment*. Retrieved from <http://www.ccg-gcc.gc.ca/e0010733>.
- Intergovernmental Panel on Climate Change. (2007). *Climate Change 2007: Synthesis Report, IPCC Fourth Assessment*. Geneva, Switzerland: IPCC.
- Melillo, J. M., T. Richmond, and G. W. Yohe. (2014). *Climate Change Impacts in the United States: The Third National Climate Assessment*. Washington, DC: U.S. Global Change Research Program.
- Minerals Management Service. (2002). *Final Environmental Impact Statement for Gulf of Mexico OCS Oil and Gas Lease Sales: 2003–2007; Central Planning Area Sales 185, 190, 194, 198, and 201; Western Planning Area Sales 187, 192, 196, and 200*. New Orleans, LA: U.S. Department of the Interior. Retrieved from <https://www.boem.gov/BOEM-Newsroom/Library/Publications/2002/2002-052-vol1.aspx>.
- Minerals Management Service. (2007a). *Programmatic Environmental Impact Statement for Alternative Energy Development and Production and Alternate Use of Facilities on the Outer Continental Shelf: Final Environmental Impact Statement*. New Orleans, LA: Gulf of Mexico OCS Region.
- Minerals Management Service. (2007b). *Gulf of Mexico OCS Oil and Gas Lease Sales: 2007–2012*. New Orleans, LA: U.S. Department of the Interior.
- National Aeronautics and Space Administration. (2018). *Why do storms generally move from west to east in the USA?* , Retrieved from <https://pmm.nasa.gov/education/content/why-do-storms-generally-move-west-east-usa>.
- National Oceanic and Atmospheric Administration. (2005). *Hurricane Katrina: A Climatological Perspective*. (Technical Report 2005-01). Asheville, NC: U.S. Department of Commerce. Retrieved from <https://www1.ncdc.noaa.gov/pub/data/techrpts/tr200501/tech-report-200501z.pdf>.
- National Oceanic and Atmospheric Administration. (2017). *National Data Buoy Center*. Retrieved from <http://www.ndbc.noaa.gov/obs.shtml>.
- National Science Foundation. (2011). *Air Quality Worsened by Paved Surfaces*. Retrieved from http://www.nsf.gov/news/news_summ.jsp?cntn_id=119606.
- Ozone Transport Commission. (2017). *What is the Ozone Transport Commission?* , Retrieved from <http://www.otcair.org/about.asp>.
- U.S. Department of Defense. (2014). *2014 Climate Change Adaptation Roadmap*. Alexandria, VA: Department of Defense.
- U.S. Department of the Navy. (1999). *Environmental Effects of RF Chaff: A Select Panel Report to the Undersecretary of Defense for Environmental Security*. Washington, DC: U.S. Department of the Navy, Naval Research Laboratory.
- U.S. Department of the Navy. (2003). *Final Environmental Impact Statement for the Introduction of the F/A-18 E/F (Super Hornet) Aircraft to the East Coast of the United States*.

- U.S. Department of the Navy. (2009). *Final Environmental Assessment Transition of E-2C Hawkeye to E-2D Advanced Hawkeye at Naval Station Norfolk, Virginia and Naval Base Ventura County Point Mugu, California*. Washington, DC.
- U.S. Department of the Navy. (2010). *United States Marine Corps F-35B East Coast Basing Environmental Impact Statement*. Norfolk, VA: Naval Facilities Engineering Command, Mid-Atlantic.
- U.S. Department of the Navy. (2013). *Navy Guidance for Compliance with the Clean Air Act General Conformity Rule*. Washington, DC: Office of the Chief of Naval Operations Energy and Environmental Readiness Division.
- U.S. Department of the Navy. (2014). *Final Supplemental Environmental Impact Statement for the Introduction of the P-8A Multi-Mission Maritime Aircraft into the U.S. Navy Fleet*. Norfolk, VA: Naval Facilities Engineering Command Atlantic.
- U.S. Environmental Protection Agency. (1972). *Mixing Heights, Wind Speeds, and Potential for Urban Air Pollution Throughout the Contiguous United States*. Research Triangle Park, NC: Environmental Protection Agency Office of Air Programs.
- U.S. Environmental Protection Agency. (2007). *The Plain English Guide to the Clean Air Act*. (EPA-456/K-07-001). Research Triangle Park, NC: U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. (2009a). *AP-42, Compilation of Air Pollutant Emissions Factors, Volume 1: Stationary Point and Area Sources*. Retrieved from <https://www.epa.gov/air-emissions-factors-and-quantification/ap-42-compilation-air-emission-factors#5thed>.
- U.S. Environmental Protection Agency. (2009b). *Endangerment and Cause or Contribute Findings for Greenhouse Gases under Section 202(a) of the Clean Air Act*. Washington, DC: U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. (2009c). *Technical support document for endangerment and cause or contribute findings for greenhouse gases under Section 202(a) of the Clean Air Act*. Washington, DC: Council on Foreign Relations.
- U.S. Environmental Protection Agency. (2010). *Exhaust and Crankcase Emission Factors for Nonroad Engine Modeling—Compression-Ignition*. (EPA-420-R-10-018). Retrieved from <https://www3.epa.gov/otaq/models/nonrdmdl/nonrdmdl2010/420r10018.pdf>.
- U.S. Environmental Protection Agency. (2016a). *Hazardous Air Pollutants*. Retrieved from <http://www.epa.gov/haps>.
- U.S. Environmental Protection Agency. (2016b). *National Ambient Air Quality Standards*. Retrieved from <https://www.epa.gov/criteria-air-pollutants/naaqs-table>.
- U.S. Environmental Protection Agency. (2016c). *Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990–2014*. Washington, DC: U.S. Environmental Protection Agency.
- U.S. Environmental Protection Agency. (2017a). *General Conformity Training Module 2.2: Emissions and Review*.
- U.S. Environmental Protection Agency. (2017b). *Nonattainment Areas for Criteria Pollutants (Green Book)*. Retrieved from <https://www.epa.gov/green-book>.
- U.S. Maritime Administration. (2016). *2015 Vessel Calls in U.S. Ports, Selected Terminals and Lightering Areas*. Retrieved from <http://www.marad.dot.gov/resources/data-statistics/#Reports>.

United Nations Framework Convention on Climate Change. (2013). *Revision of the UNFCCC Reporting Guidelines on Annual Inventories for Parties included in Annex I to the Convention (Following Incorporation of the Provisions of Decision 24/CP.19)*. (FCCC/CP/2013/10/Add.3).

This page intentionally left blank.